



Antioxidants, polyphenols, and health benefits of cherry laurel: a review

Cesarettin Alasalvar^{a*} and Sui Kiat Chang^b

^aTÜBİTAK Marmara Research Center, Food Institute, Gebze, Kocaeli, Turkey

^bKey Laboratory of Plant Resources Conservation and Sustainable Utilization, Key Laboratory of Post-Harvest Handling of Fruits, Ministry of Agriculture, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China

*Corresponding author: Cesarettin Alasalvar, TÜBİTAK Marmara Research Center, Food Institute, Gebze, Kocaeli, Turkey. Tel: +90 262 677 3200; E-mail: cesarettin.alasalvar@tubitak.gov.tr

DOI: 10.31665/JFB.2020.10226

Received: June 1, 2020; Revised received & accepted: June 24, 2020

Citation: Alasalvar, C., and Chang, S.K. (2020). Antioxidants, polyphenols, and health benefits of cherry laurel: a review. J. Food Bioact. 10: 47–52.

Abstract

Cherry laurel (dark purple or black when mature) is a popular summer fruit in the Black Sea region of Turkey. It has been gaining popularity as a valuable source of healthy fruit over the last two decades and is a good source of nutrients and polyphenols together with high antioxidant activity. This contribution discusses nutritional characteristics, antioxidants, polyphenols, and health benefits of cherry laurel. Where available, comparisons are made with other cherry fruits (Cornelian, sweet, and sour) and blueberry. Although several health benefits of this fruit and its seeds have been known as home-made and traditional medicine, *in vivo* and well-designed human clinical trials are scarce. Therefore, additional carefully-designed human clinical trials are needed to validate the health benefits of this fruit.

Keywords: Cherry laurel; Nutrients; Antioxidants; Polyphenols; Health benefits.

1. Introduction

Cherry laurel (*Laurocerasus officinalis* Roem.) belongs to the Rosacea family and is a popular fruit (dark purple or black when mature), mainly distributed and cultivated in the coasts of the Black Sea region of Turkey and is locally called “*Taflan*” or “*Karayemiş*” (Alasalvar et al., 2005; Liyana-Pathirana et al., 2006; Alasalvar, 2016). It is native to some Asian countries, the Caucasasia, Iran, some Mediterranean countries, Bulgaria, Serbia, and Western Europe (Kolayli et al., 2003; Sahan et al., 2012). Cherry laurel is mostly consumed as fresh fruit in local markets but may also be dried, pickled, and processed into syrup (known as pekmez), jam, and marmalade. Besides its use for food, both fruit and seeds of cherry laurel are well-known as traditional medicine in Turkey and have been used for many years for the treatment of stomach ulcer, digestive system complaints, bronchitis, eczemas, hemorrhoids, diuretic agent, wound healing, and hyperglycemia, among others (Baytop, 1984; Yeşilada et al., 1999; Kolayli et al.,

2003; Colak et al., 2005; Halilova and Ercisli, 2010; Turan et al., 2013; Alasalvar, 2016; Ayla et al., 2019).

This review article highlights nutritional characteristics, antioxidants, polyphenols, and health benefits of cherry laurel.

2. Nutritional characteristics of cherry laurel

The proximate compositions of cherry laurel are summarized in Table 1 (Kolayli et al., 2003; Alasalvar et al., 2005; Ozturk et al., 2017; Turkish Food Composition Database, 2020). Cherry laurel is a nutrient-dense fruit. Carbohydrate (13.96–20.23 g/100 g) is the predominant component, followed by small amounts of protein and lipid. It contains 1.61–4.02 g/100 g dietary fibre and 11.38–11.51 g/100 g sugar. Glucose is the predominant sugar, followed with fructose, in cherry laurel. With respect to minerals, cherry laurel is an excellent source of manganese (2.27–2.42 mg/100 g). The recommended dietary allowance (RDA) of manganese is 2.3 and

Table 1. Nutritional characteristics of cherry laurel (per 100 g)

Nutrient	Unit	Cherry laurel fruit ^a	References
<i>Proximate composition</i>			Alasalvar et al. (2005); Turkish Food Composition Database (2020)
Water	g	77.28–81.21	
Energy	kcal	74–76	
Protein	g	0.54–1.59	
Lipid (fat)	g	0.10–0.56	
Ash	g	0.43–0.79	
Carbohydrate	g	13.96–20.23	
Dietary fiber	g	1.61–4.02	
<i>Sugars</i>	g	11.38–11.51	Kolayli et al. (2003); Turkish Food Composition Database (2020)
Xylose	g	0.19–0.23	
Arabinose	g	0.07–0.08	
Fructose	g	4.84–5.16	
Glucose	g	5.43–5.88	
Sorbitol	g	1.51–4.80	
Sucrose	g	tr	
<i>Minerals</i>			
Calcium	mg	14.8–53	
Copper	mg	0.08–0.09	
Iron	mg	0.44–0.83	
Magnesium	mg	17.9–27.0	
Manganese	mg	2.27–2.42	
Phosphorus	mg	21–22	
Potassium	mg	157–222	
Selenium	µg	–	
Sodium	mg	1.0–5.5	
Zinc	mg	0.16–0.37	
<i>Vitamins</i>			Ozturk et al. (2017); Turkish Food Composition Database (2020)
Folate (DFE)	µg	–	
Niacin	mg	0.20–0.35	
Pantothenic acid	mg	–	
Pyridoxine (B-6)	mg	0.06–0.07	
Riboflavin	mg	0.02–0.03	
Thiamin	mg	–	
Vitamin A (RAE)	µg	tr-3.0	
Vitamin C	mg	2.00–204	
Vitamin E (ATE)	mg	–	
Vitamin K	µg	–	
<i>Carotenoids</i>			Turkish Food Composition Database (2020)
Beta-carotene	µg	tr-38	
Lutein	µg	23–35	

Abbreviations: ATE, alpha-tocopherol equivalents; DFE, dietary folate equivalents; RAE, retinol activity equivalents; tr, trace.

Table 2. Antioxidant activities of cherry laurel determined by various assays

Antioxidant assays	Unit	Cherry laurel	References
ABTS	μmol TE/g	23.21	Celep et al. (2012); Ozturk et al. (2017)
BCB	% inhibition/mL	61	
CUPRAC	mg AAE/g	23.7–46.9	
DPPH	μmol TE/g	43.54	
FRAP	μmol TE/g	14.74	
Hydrogen peroxide scavenging	% inhibition	3.5–16.3	Liyana-Pathirana et al., 2006
LDL oxidation inhibition	% inhibition	42.7–58.5	
ORAC	μmol TE/g	3,633–7,996	Alasalvar et al., 2005
Superoxide radical scavenging	% inhibition	82.3–100	Liyana-Pathirana et al., 2006

Abbreviations: AAE, ascorbic acid equivalents; ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid); BCB, beta-carotene bleaching; CUPRAC, cupric reducing antioxidant capacity; DPPH, 2,2-diphenyl-1-picrylhydrazyl; FRAP, ferric reducing antioxidant power; LDL, low-density lipoprotein; ORAC, oxygen radical absorbance capacity; TE, trolox equivalents.

1.8 mg per day for males and females, respectively (DRIs, 2001). It is also interesting to note that cherry laurel is rich in ascorbic acid (2.0–204 mg/100 g). Vitamin C, free radical scavenger, has a RDA of 90 and 75 mg per day for males and females, respectively (DRIs, 2000). Considering RDA values, consuming approximately 100 g fresh cherry laurel supplies 100% of manganese and vitamin C daily.

3. Antioxidants in cherry laurel

Antioxidant activities of cherry laurel have been reported using various assays. These include 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), beta-carotene bleaching (BCB), copper reducing antioxidant capacity (CUPRAC), 2,2-diphenyl-1-picrylhydrazyl (DPPH), ferric reducing antioxidant power (FRAP), hydrogen peroxide scavenging, low-density lipoprotein (LDL) oxidation, oxygen radical absorbance capacity (ORAC), and superoxide radical inhibitory assays (Table 2).

The antioxidant activities of cherry laurel vary according to the type of antioxidant assay used (Table 2). The antioxidant capacities of cherry laurel, determined using ABTS, DPPH, FRAP, and ORAC assays, were 23.21, 43.54, 14.74, and 3,633–7,996 μmol trolox equivalents (TE)/g, respectively (Alasalvar et al., 2005; Celep et al., 2012; Ozturk et al., 2017). Celep et al. (2012) reported that cherry laurel demonstrated similar antioxidant capacities with Cornelian cherry determined using BCB, CUPRAC, DPPH, FRAP, and superoxide radical inhibitory assays. In another study,

Capanoglu et al. (2011) reported that the antioxidant capacities of cherry laurel [28.4, 46.9, and 13.3 mmol TE/100 g dry weight (dw)] was significantly higher than sour cherry (17.8, 29.7, and 7.6 mmol TE/100 g dw), determined using ABTS, CUPRAC, and FRAP assays, respectively. However, the antioxidant capacities of both cherry laurel and sour cherry were significantly lower than Cornelian cherry. This happens since Cornelian cherry had the lowest degree of polymerization (DP) of proanthocyanidins (3.9) compared to both cherry laurel and sour cherry with high DP of proanthocyanidins (45.2 and 62.9, respectively) (Capanoglu et al., 2011). The DP is one of the most important properties in fruits and vegetables since antioxidant activity has been shown to depend on the DP (Hagerman et al., 1998; Jerez et al., 2007).

Table 3 presents contents of total anthocyanins, carotenoids, flavonoids, phenolics, and proanthocyanidins in cherry laurel (Alasalvar et al., 2005; Capanoglu et al., 2011; Celep et al., 2012; Ozturk et al., 2017). Cherry laurel contains much higher contents of total anthocyanins [124–174 mg cyanidin 3-glucoside equivalents (C3GE)/100 g fresh weight (fw)] and total phenolics [944–4,919 mg gallic acid equivalents (GAE)/100 g fw] than that of sweet cherry cultivars (0.5–29.7 mg C3GE/100 g fw and 75–407 mg GAE/100 g fw, respectively) (Chaovanalikit and Wrolstad, 2004). In addition, cherry laurel contains the same amount of total anthocyanins (40.62–378 mg C3GE/100 g fw) and higher amount of total phenolics (275–695 mg GAE/100 g fw) than that of blueberry cultivars (Rodrigues et al., 2010). In terms of total carotenoids, cherry laurel has 2-fold higher total carotenoids (254–261 μg/100 g) than that of sweet cherry (123 μg/100 g) and blueberry (112

Table 3. Contents of total polyphenols in cherry laurel

Polyphenols	Unit	Cherry laurel	References
Total anthocyanins	mg C3GE/100 g	124–174	Alasalvar et al. (2005); Capanoglu et al. (2011); Ozturk et al. (2017)
Total carotenoids	μg/100 g	254–261	
Total flavonoids	mg QE/100 g	11.76	
Total phenolics	mg FAE/100 g	454–651	
	mg GAE/100 g	944–4,919	
Total proanthocyanidins	mg EGCGE/100 g	3.40	Celep et al. (2012)

Abbreviations: C3GE, cyanidin 3-glucoside equivalents; EGCGE, epigallocatechin gallate equivalents; FAE, ferulic acid equivalents; GAE, gallic acid equivalents; QE, quercetin equivalents.

Table 4. Reported polyphenols in cherry laurel

Types of polyphenolic compounds		Unit	Content	References
Anthocyanins	Cyanidin-3-glucoside	mg/100 g fw	4.6	Capanoglu et al. (2011); Karahalil and Şahin (2011); Bayrambaş et al. (2019);
	Cyanidin-3-rutinoside		6.6	
	Keracyanin chloride		2.4–14.97	
	Pelargonidin-3-glucoside		11.0	
Flavan-3-ols	(-)-Catechin		0.02–23.3	
	(-)-Catechin gallate		0.8–4.3	
	(-)-Epigallocatechin		1.7–27.2	
	(-)-Epigallocatechin gallate		0.4–41	
	Epicatechin		0.3–17.2	
	(-)-Gallocatechin gallate		1.4–6.4	
Flavonols	Kaempferol		0.5–3.8	
	Quercetin-3-glucoside		0.97–2.8	
	Rutin		nd–0.3	
Phenolic acids	Caffeic acid		64.6–74.8	Alasalvar et al. (2005); Capanoglu et al. (2011); Karahalil and Şahin (2011); Karabegović et al. (2014); Ozturk et al. (2017); Bayrambaş et al. (2019)
	Chlorogenic acid		103–160	
	<i>o</i> -Coumaric acid		nd–2.24	
	<i>p</i> -Coumaric acid		11.5–49.7	
	Ferulic acid		nd–6.2	
	Gallic acid		0.2–37.6	
	<i>p</i> -Hydroxybenzoic acid		2.3–3.0	
	Protocatechuic acid		nd–1.12	
	Protocatechuic acid ethyl ester		0.04–19.0	
	Syringic acid		1.0–56.2	
	Vanillic acid		1.5–12.4	
Procyanidins	Procyanidin B2		14.4–52.4	

Abbreviations: fw, fresh weight; nd, not detected.

µg/100 g) (USDA, 2019). Thus, cherry laurel may be considered as a rich source of polyphenols.

4. Polyphenols in cherry laurel

To the best of our knowledge, limited polyphenols (anthocyanins, flavan-3-ols, flavonols, phenolic acids, and procyanidins) have been reported in cherry laurel (Table 4). Phenolic acids (11 in total) are the main polyphenols reported in cherry laurel. Chlorogenic acid is the most abundant phenolic acid, followed by caffeic, *p*-coumaric, syringic, and gallic acids. Besides, *o*-coumaric, ferulic, *p*-hydroxybenzoic, protocatechuic, protocatechuic acid ethyl ester, and vanillic acids are also present (Alasalvar et al., 2005; Karahalil & Şahin, 2011; Karabegović et al., 2014; Ozturk et al., 2017; Bayrambaş et al., 2019). Ayaz et al. (1997) reported the presence of protocatechuic, *p*-hydroxybenzoic, vanillic, caffeic, and *p*-coumaric acids in several cultivated and wild cherry laurel varieties where vanillic acid was the most abundant phenolic acid.

Four anthocyanins have been reported in cherry laurel (Table 4). Keracyanin chloride was the most abundant anthocyanins, as

reported in a recent study (Bayrambaş et al., 2019). Besides, cyanidin-3-glucoside, cyanidin-3-rutinoside, and pelargonidin-3-glucoside were also detected in cherry laurel (Karahalil and Şahin, 2011).

Capanoglu et al. (2011) reported that the total flavan-3-ol content in cherry laurel was 1,089 mg/100 g dw where (-)-epigallocatechin gallate and (-)-epigallocatechin were the most abundant flavan-3-ols. Besides, kaempferol was the most abundant flavonols detected in cherry laurel, followed by quercetin-3-glucoside and rutin. Cherry laurel also contains procyanidin B2 (14.4–52.5 mg/100 g fw) (Bayrambaş et al., 2019). Figure 1 shows chemical structures of the phenols and polyphenols present in cherry laurel.

5. Health benefits of cherry laurel

Since cherry laurel has high contents of phenolic acids, anthocyanins, and flavan-3-ols, it is able to provide health benefits beyond basic nutrition (Karahalil and Şahin, 2011; Bayrambaş et al., 2019). Cherry laurel renders beneficial effects on human health such as




Figure 1. Chemical structures of the phenols and polyphenols present in cherry laurel.

neuroprotective, anti-diabetic, dimethoate-induced liver damage, antioxidative stress, anti-microbial, wound healing, and cytotoxic activities (Orhan and Akkol, 2011; Orhan et al., 2015; Demir et al., 2017; Eken et al., 2017; Ayla et al., 2019; Bayrambaş et al., 2019). These health effects of cherry laurel are reviewed below.

Orhan and Akkol (2011) evaluated the neuroprotective potential of cherry laurel extract. Results demonstrated that dichloromethane extract of cherry laurel (100 and 200 µg/mL) showed 16.3 and 19% inhibitory activity on acetylcholinesterase, respectively, but had no effect on the activity of butyrylcholinesterase enzyme. These enzymes play important roles in the pathogenesis of Alzheimer's disease (Orhan and Akkol, 2011). More research should be conducted to ascertain the neuroprotective potential of cherry laurel.

Orhan et al. (2015) determined the effect of ethanolic extract of pitted and non-pitted cherry laurel on blood glucose levels in healthy glycemic, glucose-loaded, and streptozotocin-induced diabetic rats. Results demonstrated that pitted extract of cherry laurel (500 and 1,000 mg/kg) showed a notable hypoglycemic effect in healthy male rats. However, pitted cherry laurel extract (1,000 and 2,000 mg/kg) had weak inhibitory effect on blood glucose levels in glucose-loaded rats. All cherry laurel extracts had no effect on blood glucose levels in streptozotocin-induced diabetic rats (Orhan et al., 2015).

More recently, Eken et al. (2017) evaluated the protective role of methanolic extract of cherry laurel against dimethoate-induced liver damage and oxidative stress *in vivo*. Their results showed that pre- and post-treatment with cherry laurel extract significantly reduced the activity of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) as well as total bilirubin level compared to the control group. Besides, administration of cherry laurel extract

also normalized total antioxidant status by increasing the enzymatic activities of superoxide dismutase, catalase and glutathione peroxidase, reducing the malondialdehyde level, and DNA damage significantly (Eken et al., 2017). Supplementation with cherry laurel extract protected liver against hepatotoxicity by improving the structural integrity of hepatocyte cell membrane and regeneration of damaged liver cells, thus maintaining normal hepatic physiology. The liver-protective effects of cherry laurel extract were comparable to the positive control, vitamin C (Eken et al., 2017).

The antimicrobial effects of methanolic extract of cherry laurel against five microbial strains including Gram-positive bacteria (*Bacillus cereus* and *Staphylococcus aureus*) and Gram-negative bacteria (*Escherichia coli*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa*) was determined by Ayla et al. (2019), using agar diffusion method. Cherry laurel extract exhibited potential inhibitory activities against growth of the five tested human pathogenic microbes where there were no significant differences between the antimicrobial effects between Gram-positive and Gram-negative bacteria. In addition, cherry laurel extract was also investigated for wound healing effects in excisional wound mice model for ten days. Results demonstrated that topical application of cherry laurel extract on wound area accelerated wound contraction rate and enhanced epithelialization and angiogenesis in the wound healing compared to the control group (Ayla et al., 2019). The authors postulated that the antioxidant and antimicrobial effects of cherry laurel contributed to the wound healing effects.

Finally, Bayrambaş et al. (2019) investigated the cytotoxic effect of methanolic extract of three cherry laurel varieties on HCT 116 human colon cancer cell line. Treatment with 61K04 variety prevented the growth of cancer cells (55% apoptosis), followed by 61K05 (27% apoptosis), and 55K06 (36.9% apoptosis) varie-

ties. In another study, Demir et al. (2017) reported that dimethyl sulphoxide extract of cherry laurel exhibited potential cytotoxic effect on human colon cancer cells (WiDr) and human lung cancer cells (A549) *in vitro*. However, cherry laurel extract did not show any effect on the cell viability of human prostate cancer cells (PC-3), hepatocarcinoma cells (HepG2), cervix adenocarcinoma cells (HeLa), and breast adenocarcinoma cells (MCF-7). These findings should be validated in animal models to provide better indication of the use of cherry laurel extract in preventing severity of various types of cancer.

6. Conclusion

The available data to date reveal that cherry laurel is rich in antioxidant phenolics and possesses strong radical scavenging activity. It is a good source of nutrients and polyphenols (mainly phenolic acids, anthocyanins, and flavan-3-ols). Limited data exist in the literature on the detailed polyphenol profiles of cherry laurel. Further research is warranted to identify other polyphenols present in cherry laurel such as anthocyanins, ellagitannins, and proanthocyanidins as well as to addressing its health benefits using well-designed human clinical trials.

References

- Alasalvar, C. (2016). Cherry laurel syrup (Pekmez). In: Shahidi, F., and Alasalvar, C. (Ed.). Handbook of functional beverages and human health. CRC Press, Taylor & Francis Group, Boca Raton, FL, pp. 187–192.
- Alasalvar, C., Al-Farsi, M., and Shahidi, F. (2005). Compositional characteristics and antioxidant components of cherry laurel varieties and pekmez. *J. Food Sci.* 70: S47–S52.
- Ayaz, F.A., Kadioğlu, A., Reunanen, M., and Var, M. (1997). Phenolic acid and fatty acid composition in the fruits of *Laurocerasus officinalis* Roem. and its cultivars. *J. Food Compos. Anal.* 10: 350–357.
- Ayla, S., Okur, M.E., Günel, M.Y., Özdemir, E.M., Çiçek Polat, D., Yoltaş, A., Biçeroglu, Ö., and Karahüseyinoğlu, S. (2019). Wound healing effects of methanol extract of *Laurocerasus officinalis* Roem. *Biotech. Histochem.* 94: 180–188.
- Bayrambaş, K., Çakır, B., and Gülseren, İ. (2019). Influence of phenolic profile on the RP-HPLC detection and anti-carcinogenic potential of cherry laurel extracts from Black Sea Region-Turkey. *Microchem. J.* 149: 103963.
- Baytop, T. (1984). Therapy with medicinal plants in Turkey (past and present). Istanbul University Publication No. 3255, Istanbul, Turkey.
- Capanoglu, E., Boyacioglu, D., de Vos, R.C., Hall, R.D., and Beekwilder, J. (2011). Procyanidins in fruit from sour cherry (*Prunus cerasus*) differ strongly in chain length from those in laurel cherry (*Prunus laurocerasus*) and Cornelian cherry (*Cornus mas*). *J. Berry Res.* 1: 137–146.
- Celep, E., Aydın, A., and Yesilada, E. (2012). A comparative study on the *in vitro* antioxidant potentials of three edible fruits: Cornelian cherry, Japanese persimmon and cherry laurel. *Food Chem. Toxicol.* 50: 3329–3335.
- Chaovanalikit, A., and Wrolstad, R.E. (2004). Total anthocyanins and total phenolics of fresh and processed cherries and their antioxidant properties. *J. Food Sci.* 69: FCT67–FCT72.
- Colak, A., Özen, A., Dincer, B., Güner, S., and Ayaz, F.A. (2005). Diphenolases from two cultivars of cherry laurel (*Laurocerasus officinalis* Roem.) fruits at an early stage of maturation. *Food Chem.* 90: 801–807.
- Demir, S., Turan, I., Demir, F., Demir, E.A., and Aliyazicioglu, Y. (2017). Cytotoxic effect of *Laurocerasus officinalis* extract on human cancer cell lines. *Marmara Pharm. J.* 21: 121–126.
- DRIs. (2000). DRIs. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. The National Academies Press, Washington, DC.
- DRIs. (2001). DRIs. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. The National Academies Press, Washington, DC.
- Eken, A., Endirlik, B.Ü., Bakir, E., Baldemir, A., Yay, A.H., and Canturk, F. (2017). Effect of *Laurocerasus officinalis* Roem. (cherry laurel) fruit on dimethoate induced hepatotoxicity in rats. *Kafkas Üniv. Vet. Fak. Derg.* 23: 779–787.
- Hagerman, A.E., Riedl, K.M., Jones, G.A., Sovik, K.N., Ritchard, N.T., Hartzfeld, P.W., and Riechel, T.L. (1998). High molecular weight plant polyphenolics (tannins) as biological antioxidants. *J. Agric. Food Chem.* 46: 1887–1892.
- Halilova, H., and Ercisli, S. (2010). Several physico-chemical characteristics of cherry laurel (*Laurocerasus officinalis* Roem.) fruits. *Biotechnol. Biotechnol. Eq.* 24: 1970–1973.
- Jerez, M., Selga, A., Sineiro, J., Torres, J.L., and Nunez, M.J. (2007). A comparison between dark extracts from *Pinus pinaster* and *Pinus radiata*: Antioxidant activity and procyanidins composition. *Food Chem.* 100: 439–444.
- Karabegović, I.T., Stojičević, S.S., Veličković, D.T., Todorović, Z.B., Nikolić, N.Č., and Lazić, M.L. (2014). The effect of different extraction techniques on the composition and antioxidant activity of cherry laurel (*Prunus laurocerasus*) leaf and fruit extracts. *Ind. Crops Prod.* 54: 142–148.
- Karahalil, F.Y., and Şahin, H. (2011). Phenolic composition and antioxidant capacity of cherry laurel (*Laurocerasus officinalis* Roem.) sampled from Trabzon region, Turkey. *Afr. J. Biotechnol.* 10: 16293–16299.
- Kolaylı, S., Küçük, M., Duran, C., Candan, F., and Dinçer, B. (2003). Chemical and antioxidant properties of *Laurocerasus officinalis* Roem. (Cherry laurel) fruit grown in the Black Sea region. *J. Agric. Food Chem.* 51: 7489–7494.
- Liyana-Pathirana, C.M., Shahidi, F., and Alasalvar, C. (2006). Antioxidant activity of cherry laurel (*Laurocerasus officinalis* Roem.) and its concentrated juice. *Food Chem.* 99: 121–128.
- Orhan, I.E., and Akkol, E.K. (2011). Estimation of neuroprotective effects of *Laurocerasus officinalis* Roem. (cherry laurel) by *in vitro* methods. *Food Res. Int.* 44: 818–822.
- Orhan, N., Damlaçi, T., Baykal, T., Özek, T., and Aslan, M. (2015). Hypoglycaemic effect of seed and fruit extracts of laurel cherry in different experimental models and chemical characterization of the seed extract. *Rec. Nat. Prod.* 9: 379.
- Ozturk, B., Celik, S.M., Karakaya, M., Karakaya, O., Islam, A., and Yarılgac, T. (2017). Storage temperature affects phenolic content, antioxidant activity and fruit quality parameters of cherry laurel (*Prunus laurocerasus* L.). *J. Food Process. Preserv.* 41: e12774.
- Rodrigues, E., Poerner, N., Rockenbach, I.I., Gonzaga, L.V., Mendes, C.R., and Fett, R. (2010). Phenolic compounds and antioxidant activity of blueberry cultivars grown in Brazil. *Ciênc. Tecnol. Aliment., Campinas* 31: 911–917.
- Sahan, Y., Cansev, A., Celik, G., and Cinar, A. (2012). Determination of various chemical properties, total phenolic contents, antioxidant capacity and organic acids in *Laurocerasus officinalis* fruits. *Acta Hort.* 939: 359–366.
- Turan, M.I., Turkoglu, M., Dundar, C., Celik, N., and Suleyman, H. (2013). Investigating the effect of *Prunus laurocerasus* fruit extract in type II diabetes induced rats. *Int. J. Pharmacol.* 9: 373–378.
- Turkish Food Composition Database. (2020). Turkish Food Composition Database. Cherry laurel. Published online at <http://www.turkomp.gov.tr/food-cherry-laurel-570> (accessed June 20, 2020).
- USDA. (2019). USDA FoodData central for SR legacy. Published online at <https://fdc.nal.usda.gov> (accessed June 20, 2020).
- Yeşilada, E., Sezik, E., Honda, G., Takaishi, Y., Takeda, Y., and Tanaka, T. (1999). Traditional medicine in Turkey IX: folk medicine in north-west Anatolia. *J. Ethnopharmacol.* 64: 195–210.