

5-Demethylnobiletin more potently inhibits colon cancer cell growth than nobiletin *in vitro* and *in vivo*

Yi-Siou Chiou^a, Yu-Nu Zheng^b, Mei-Ling Tsai^b, Ching-Shu Lai^b,
Chi-Tang Ho^c and Min-Hsiung Pan^{a,d,e,*}

^aInstitute of Food Science and Technology, National Taiwan University, Taipei 10617, Taiwan

^bDepartment of Seafood Science, National Kaohsiung Marine University, Kaohsiung 811, Taiwan

^cDepartment of Food Science, Rutgers University, New Brunswick, NJ 08901, USA

^dDepartment of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan

^eDepartment of Health and Nutrition Biotechnology, Asia University, Taichung, Taiwan

*Corresponding author: Min-Hsiung Pan, Institute of Food Science and Technology, National Taiwan University, No. 1, Section 4, Roosevelt Road, Taipei 10617, Taiwan. Tel: +886 2 33664133; Fax: +886-2-33661771; E-mail: mhpan@ntu.edu.tw

DOI: 10.31665/JFB.2018.2143

Received: January 15, 2018; Revised received & accepted: February 19, 2018

Citation: Chiou, Y.-S., Zheng, Y.-N., Tsai, M.-L., Lai, C.-S., Ho, C.-T., and Pan, M.-H. (2018). 5-Demethylnobiletin more potently inhibits colon cancer cell growth than nobiletin *in vitro* and *in vivo*. J. Food Bioact. 2: 91–97.

Abstract

Nobiletin (NOB) and 5-demethylnobiletin (DMNB) are unique polymethoxyflavones (PMFs) found in citrus peel that exhibit anti-tumoral action in several cancer cell models. The differences between NOB and DMNB with respect to their anti-proliferative potencies and underlying molecular mechanism were compared in this contribution. The results of the cell viability assay suggested that DMNB resulted in more enhanced growth inhibitory effects than NOB in human colon cancer cell lines (HCT-116, HT-29 and COLO 205). Flow cytometry data found that DMNB inhibited proliferation in COLO 205 cells by predominantly inducing apoptosis. A xenograft mouse model further demonstrated that DMNB exhibited more preferential anti-colon cancer effects than NOB via its ability to induce p53-regulated cell death signaling (apoptosis and autophagy) and inhibit key cellular markers associated with inflammation and angiogenesis. Taken together, our findings provide evidence for the first time that natural bioactive DMNB might serve as a promising polymethoxyflavone for chemoprevention of colorectal cancer.

Keywords: 5-Demethylnobiletin; Nobiletin; Colorectal cancer; p53; Apoptosis; Autophagy.

1. Introduction

Colorectal cancer (CRC) is the third most common malignancy and has been ranked the second driving reason for death from cancer among both men and women (Jemal et al., 2010). Although the dissemination of screening tests has led to a decline in CRC incidence and mortality over the last two decades (Simon, 2016), changes in some modifiable factors such as eating habits and lifestyle patterns have been positively or negatively associated with the risk of CRC development (Tuan and Chen, 2016). Furthermore, the current treatment modalities including surgery, chemotherapy and radiotherapy for colon cancer are limited due to resist-

ance, undesirable side effects and complications, finally leading to therapeutic failure (Denlinger and Barsevick, 2009; Hu et al., 2016). Therefore, there is an urgent need for effective approaches and safer medicines to overcome this public health challenge and improve the quality of life of patients.

Recently, epidemiological and experimental evidence has demonstrated that a diet containing high amounts of fat, red meat and refined sugar appear to increase CRC risk appreciably. Conversely, consumption of dietary phytochemicals from vegetables, fruits and spices has the potential to prevent CRC (Li et al., 2015). Therefore, the use of dietary bioactive compounds either alone or in combination to prevent or treat cancer has received much attention due to their safety record and green status. Polymethoxyflavones (PMFs)

belong to the family of unique flavonoids that specifically exist in citrus peels such as sweet oranges (*Citrus sinensis*), mandarin oranges (*Citrus reticulata*) and miaray mandarin (*Citrus miaray*) (Li et al., 2006; Uckoo et al., 2015). The compound 5-demethylnobiletin (DMNB) is an autohydrolysis product of nobiletin (NOB) during storage (Zheng et al., 2013), and both are naturally occurring bioactive constituents of PMFs (Wang et al., 2007). Current studies have shown that NOB enhances the effects of cancer cell growth inhibition in various cancer cell lines such as those of gastric, colorectal, lung and breast (Luo et al., 2008; Morley et al., 2007; Moon and Cho, 2016; Moon et al., 2013; Yoshimizu et al., 2004; Chen et al., 2014; Sp et al., 2017). DMNB has been reported to have rather strong pharmacological properties including anti-atherogenic (Yen et al., 2011), neurotrophic (Chiu et al., 2013), anti-obesity (Tung et al., 2016), anti-lung cancer (Chen et al., 2015; Song et al., 2016; Song et al., 2017) efficacies. Thus, it has been implied that both NOB and DMNB may act as effective and diverse cytostatic anticancer agents.

The NOB and its colonic metabolites have excellent chemopreventive activity in a colitis-associated colon carcinogenesis mouse model as well as in human colon cancer cell models (Morley et al., 2007; Kawabata et al., 2005; Wu et al., 2017; Wu et al., 2015). However, no information is available on the effects of DMNB on colon cancer cell growth inhibition. More importantly, we have previously found that dietary administration of hydroxylated PMFs produces a stronger chemopreventive effect in azoxymethane (AOM)-induced colonic tumorigenesis (Lai et al., 2011). However, the effects of bioactive hydroxylated PMFs on colon cancer cell growth inhibition have not been systematically elucidated. Therefore, the present study aimed to compare both *in vitro* and *in vivo* anti-proliferative effects of NOB and DMNB, and elucidate the underlying molecular mechanisms.

To the best of our knowledge, this work is the first study to demonstrate that DMNB exerts higher anti-colon cancer efficacies than NOB, which is related to its proapoptotic, pro-death autophagy, anti-inflammatory, and anti-angiogenic activities. These findings support the use of DMNB in colorectal cancer chemoprevention and chemotherapy.

2. Materials and methods

2.1. Chemicals and reagents

The compound NOB was purchased from Quality Phytochemicals LLC (Edison, NJ, USA), while DMNB was synthesized and obtained as a pale yellow crystal solid through acid hydrolysis from NOB as described previously (Zheng et al., 2013). Antibodies against poly-ADP-ribose polymerases (PARP), p53, vascular endothelial growth factor (VEGF) and β -actin were purchased from Cell Signaling Technology (Beverly, MA, USA). Anti-cyclooxygenase-2 (COX-2) was purchased from BD Transduction Laboratories (Lexington, KY, USA). Anti-light chain 3 (LC3) was purchased from MBL International (San Diego, CA, USA). All other chemicals used in this study were of the highest purity and were commercially available.

2.2. Cell culture

Human colon cancer cells were purchased from the American Type Culture Collection (Rockville, MD, USA) and maintained in 90% RPMI-1640 medium and 10% fetal bovine serum (Gibco BRL, Grand Island, NY, USA), supplemented with 2 mM glutamine

(Gibco BRL), 1% penicillin/streptomycin (100 U of penicillin/mL and 100 μ g streptomycin/mL) in a humidified atmosphere containing 5% CO₂ at 37 °C.

2.3. Cell viability assay

Cell viability was determined for indicated compounds (NOB and DMNB) based on the classical MTT assay. Briefly, human colon cancer HCT-116, HT-29 and COLO 205 cells were plated at a density of 1×10^5 cells/mL into 96-well plates. After overnight growth, cells were treated with a series of concentrations of NOB and DMNB for 24 h. The final concentrations of dimethyl sulfoxide (DMSO) in the culture medium were <0.05%. The viability percentage was calculated based on the percentage of unstained cells, as described previously (Pan et al., 2007).

2.4. Apoptosis detection

COLO 205 cells 1×10^5 cells/mL were cultured in 24-well plates and incubated overnight. The cells were treated with various concentrations (5, 10, and 20 μ M) of NOB and DMNB for 24 h. The cells were then harvested, washed with PBS, resuspended in 200 μ L of phosphate-buffered saline (PBS), and fixed in 800 μ L of iced 100% ethanol at -20 °C. After being left to stand overnight, the cell pellets were collected by centrifugation, resuspended in 500 μ L of hypotonic buffer (0.5% Triton X-100 in PBS) and 0.5 μ g/mL RNase), and incubated at 37 °C for 15 min. Subsequently, propidium iodide (PI) solution (10 μ g/mL) was added, and the mixture was allowed to stand at 37 °C for 15 min. Fluorescence emitted from the PI-DNA complex was quantified after excitation of the fluorescent dye by FACSscan cytometry (Becton Dickinson, San Jose, CA, USA).

2.5. Xenograft model establishment

Five-week-old male BALB/cAn.Cg-Foxnl^{nu}/Crlnarl mice were purchased from the BioLASCO Experimental Animal Center (Taiwan Co., Ltd, Taipei, Taiwan). All animal experimental protocols used in this study were approved by Institutional Animal Care and Use Committee of the National Kaohsiung Marine University (IAC UC, NKMU) and maintained in pathogen-free sterile isolators according to institutional guidelines and all food, water, caging and bedding were sterilized before use. To assess the efficacy of NOB and DMNB against tumor growth, COLO 205 cells (5×10^6 cell/100 μ L) were subcutaneously injected into the ventral flank of each nude mouse. When the diameter of all xenograft tumors reached 100 mm³, the model was considered successfully established and treatment was initiated.

2.6. Animal grouping and treatment

Tumor-bearing mice were randomized into four groups (n = 6) and injected intraperitoneally (*ip*) daily with DMSO solvent, NOB 100 mg/kg and DMNB 50 and 100 mg/kg, respectively, for 21 days. At the beginning of treatment (day 0), the body weight and tumor volume were measured once every three days. The maximal diameter of the tumor (a) and the short diameter (b) were calculated with a digital caliper. The tumor volume was estimated by using the formula: volume = $\frac{1}{2} \times a \times b^2$. On day 21, all nude mice were sacrificed and the tumors were excised and then weighed using an electronic balance. During the administration, the activity and diet were also observed.




Figure 1. Effect of NOB and DMNB on the growth of HCT-116, HT-29 and COLO 205 cells. (A) Chemical structures of NOB and DMNB. (B) Cell well treated with 0, 1, 2.5, 5, 10, 20 and 40 μM of NOB or DMNB for 24 h. Cell viability was determined by MTT assay. Data are expressed as mean \pm SD. When there were significant differences among control, NOB and DMNB groups, they were further analyzed by one-way ANOVA and Duncan's Multiple Range Test, with results being indicated by the letters a, b, c, and d.

2.7. Western blotting

Following experimental treatments, all tumors were harvested and lysed with lysis buffer [10% glycerol, 1% Triton X-100, 1 mM Na₃VO₄, 1 mM ethylene glycol tetraacetic acid (EGTA), 10 mM NaF, 1 mM Na₄P₂O₇, 20 mM Tris buffer (pH7.9), 100 μM β -glycerophosphate, 137 mM NaCl, and 5 mM ethylenediamine-tetraacetic acid (EDTA)] containing 1 Protease Inhibitor Cocktail Tablet (Roche, Indianapolis, IN, USA) on ice for 1 h, followed by centrifugation at $17,500 \times g$ for 30 min at 4 °C. Total proteins were extracted as previously described (Chiou et al., 2013). For immunoblot analysis, equal amounts of lysates (50 μg) were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and electrically transferred to a polyvinylidene difluoride (PVDF) membrane (Merck Millipore, Darmstadt, Germany). After blocking, the membrane was incubated overnight with specific primary antibodies at 4 °C, washed three times with PBS-tween 20 (PBST), and then incubated with horseradish peroxidase (HRP)-conjugated secondary antibodies at room temperature for 1h. The membrane was then exposed to X-ray film (FUJIFILM, Super RX, Düsseldorf, Germany) (Eastman Kodak Co., Rochester, NY, USA) and the protein bands were visualized using ECL chemiluminescence reagents (Amersham, Arlington Heights, IL, USA).

2.8. Statistical analysis

All data were repeated in a minimum of three separate experiments,

and the data were presented as the mean \pm standard deviation (SD) or standard error (SE). SigmaPlot 10.0 software was used for illustration and analysis. *P* values were calculated using Student's *t*-test or one-way ANOVA followed by Duncan's multiple-range comparison. A *P* value < 0.05 was considered statistically significant.

3. Results and discussion

3.1. DMNB showed stronger cytostatic effects than NOB on colon cancer cells

Despite significant interest in the health-promoting effects of PMFs such as tangeretin (TAN) and NOB, hydroxylated PMFs have been demonstrated to exhibit more potent biological activities than their permethoxylated counterpart PMFs (Sergeev et al., 2007; Charoensinphon et al., 2013; Ma et al., 2014; Kimura et al., 2013). Our previous study has also shown that hydroxylated PMFs cause much stronger growth inhibitory effects on colon cancer cells (Qiu et al., 2010). Here, we therefore investigated the structure-activity relationships between NOB and DMNB on three human colon cancer cell lines: HCT-116, HT-29 and COLO205. The structures of both PMFs is illustrated in Fig. 1A. The MTT assay was performed to examine the influence of NOB and DMNB on cell viability. Fig. 1B illustrates that DMNB significantly reduced cell survival on three human colon cancer cell lines at high concentrations (5–40 μM) in the following order: COLO205>HCT-116>HT-29, but NOB




Figure 2. Apoptotic effects of NOB and DMNB on COLO 205 cells. (A) Cell morphology of the control group, 20 μM NOB group and 20 μM DMNB group. Cell well treated with 5, 10 and 20 μM of NOB or DMNB for 24 h. (B) The apoptosis ratio (%) was detected by flow cytometry and quantification of sub-G1 phase in COLO 205 cells. Data were expressed as mean \pm SD. When there were significant differences among control, NOB and DMNB groups, they were further analyzed by one-way ANOVA and Duncan's Multiple Range Test, and results are indicated by the letters a, b, and c. (C) The expression of PARP protein was detected by using a Western blot with a specific antibody.

showed little effect at the same concentrations. However, at 40 μM , NOB had a markedly pronounced cytotoxicity, but this effect was lower than that of DMNB treatment (Fig. 1B). On the other hand, DMNB may be more effective than NOB against colon cancer, indicating the pivotal role of the 5'-hydroxyl group on the A-ring moiety of natural PMFs for enhanced anti-proliferative activity.

3.2. DMNB caused cell death through induction of pro-apoptotic signaling

Programmed cell death (PCD), referred to as apoptosis, is a homeostatic mechanism to maintain cellular balance and genome integrity via removing aging or unhealthy cells (Ouyang et al., 2012). It is generally believed that inappropriate apoptosis or uncontrolled cell proliferation may be an important factor in cancer development. Furthermore, evasion of apoptosis is a fundamentally important factor to serve both tumor progression and chemo-resistance in CRC (Abraha and Ketema, 2016). Thus, targeting the apoptotic pathway is considered vital in colorectal cancer therapies. Previously, DMNB has been reported to exert an anti-lung cancer effect with its induction of tumor cell cycle arrest, apoptosis and autophagy (Song et al., 2017; Chen et al., 2015; Charoensinphon et al., 2013). By applying the same rationale, we wanted to determine whether the DMNB-induced growth inhibition was caused by the induction of apoptosis.

As Figure 2A illustrates, when COLO 205 cells were treated

with DMNB at a high concentration (20 μM) for 24 h, the characteristics of apoptosis such as shrunken, apoptotic bodies and cell detachment were clearly visible. However, these morphological changes were absent from the control and NOB-treated cells (Figure 2A). Flow cytometry analysis demonstrated a significant apoptosis ratio (~26%) induced by DMNB, while NOB treatment at any concentration was not observed (Figure 2B). In addition, we also adopted immunoblotting to confirm that DMNB significantly induced apoptosis in a dose-dependent manner (Figure 2C), with application of 10 and 20 μM DMNB resulting in cleavage of PARP (cPARP), which is the hallmark of apoptosis. These data indicate that pro-apoptotic activity is critical for enhancing DMNB-induced cell death up to a level of 50%.

3.3. DMNB had a higher potential than NOB as a chemotherapeutic agent against colon cancer

Clinical chemotherapeutic drugs show many toxic side effects, which places significant limitations on their clinical application. In contrast, healthy dietary patterns used for cancer risk reduction have been approved. The naturally occurring constituents of the human diet have been intensively valued in developing new medicines and functional foods for the cancer chemoprevention field. To further confirm that DNMB-induced anticancer activities are better than NOB, we established a colorectal cancer xenograft mouse model and determined its effects on tumor growth and the




Figure 3. DMNB effectively inhibits tumor growth of COLO 205 tumor xenografts in nude mice. (A) The change of body weight and (B) tumor volume from each group during the experiment. (C) Representative xenograftic tumors from the vehicle-treated positive, NOB (50 mg/kg) and DMNB (50 and 100 mg/kg)-treated groups (day 21). (D) Final tumor weights were also measured and plotted. Data were expressed as mean ± SE. When there were significant differences among positive, NOB and DMNB groups, they were further analyzed by one-way ANOVA and Duncan’s Multiple Range Test, and results are indicated by the letters a and b.

related molecular mechanisms. Over the treatment period, the body weight of nude mice in each group did not differ and no signs of toxicity were observed, suggesting that long-term treatment of both PMFs did not cause any noticeable side effects (Figure 3A). In contrast, the tumor growth inhibition was most evident in mice treated with 50 and 100 mg/kg of DMNB via *i.p* injection. As shown in Figure 3B, the tumor volumes in the 50 and 100 mg/kg DNMB groups were stabilized in a steady state starting from Day 12, and were much slower than the positive or NOB-treated groups. In addition, the eventual tumor size and tumor weight of the DMNB-treated group were also much smaller than the control ($P < 0.05$) at the end of the experiment (Figure 3C and 3D).

To validate our *in vitro* results, we also detected apoptosis markers. As shown in Figure 4A, after treatment with 50 and 100 mg/kg of DMNB, the ratio of cleaved-PARP was increased in tumor tissues. Previously, DMNB-induced autophagy was found to be associated with mitigated apoptosis (Chen et al., 2015), which exemplifies the importance of autophagy signaling in cell death. Inconsistent with the results of the previous study, the expression of LC3, which is a key regulator involved in the formation of the autophagosome, was significantly increased (Figure 4A), suggesting that DMNB-induced cell death was closely correlated with the pro-death function of autophagy, but predominantly through apoptosis.

The p53 signaling pathways are considered to be an interconnected network for regulation of apoptosis and autophagy (Nikolopoulou et al., 2013). Owing to the multiplicity of its function, mutation or dysregulation of p53 is tightly associated with tumor

progression. In order to characterize the involvement of p53 signaling in DMNB-induced cell death, we detected the p53 protein level. As shown in Figure 4B, promising results were obtained from western blotting and confirmed that p53 up-regulation by DNMB could actually provoke the pro-apoptotic and pro-death autophagy-targeted genes, a novel finding.

To further explore how DNMB exerted its antitumor effects by interfering with pro-inflammatory and pro-angiogenic pathways, we also detected the expression of COX-2 and VEGF after DMNB treatment. As shown in Fig. 4B, tumor-related increases of COX-2 and VEGF expression were markedly impeded in the presence of DNMB, and these effects were also stronger than that of NOB. The results suggest that the anti-proliferative effect of DNMB was accompanied by inducing pro-death signaling (apoptosis and autophagy), anti-inflammation and anti-angiogenesis (Figure 5). Thus, these findings demonstrate that DNMB had higher potential than NOB as a useful approach for colon cancer therapeutics.

4. Conclusion

This is the first report showing that the naturally hydroxylated PMF, DNMB, holds much higher potential promise for colon cancer treatment than NOB in both *in vitro* and *in vivo* studies, an effect that is significantly mediated by its pro-apoptotic and pro-death autophagy properties. More importantly, we found for the first time that specifically increasing p53 by DNMB might be in-




Figure 4. Effects of DMNB on pro-death, inflammation and angiogenesis signaling pathways in tumor tissue. Mice were treated as described in Materials and Methods. At the end of the experiment, tumor tissues were excised and protein lysates of homogenized tissue were measured for PARP, LC3, p53, COX-2 and VEGF expression by Western blot analysis. Quantification of target protein expression was normalized to β-actin using a densitometer by Image J software.




Figure 5. Schematic diagram showing the mechanisms underlying the anti-colon cancer effects of DMNB. These anti-proliferative potencies were accompanied by inducing p53-regulated cell death signaling (apoptosis and autophagy) and cooperating inhibition of key cellular proteins (COX-2 and VEGF) associated with inflammation and angiogenesis. Abbreviation: DMNB, 5-demethylnobiletin; cPARP, cleaved poly-ADP-ribose polymerases; LC3II, light chain 3 II; COX-2, cyclooxygenase-2; and VEGF, vascular endothelial growth factor.

involved in cell death signaling. This strongly implies that DMNB may have potential as an ideal therapeutic agent, particularly for colorectal cancer patients with wild-type p53 expression. Furthermore, results presented here also reveal that DMNB produced anti-inflammatory and anti-angiogenic efficacies which play a synergistic role in further enhancing the growth inhibition induced by DMNB. Overall, these findings provide a powerful theoretical and experimental basis for developing DMNB as a more effective and safer agent and functional food for colon cancer prevention and therapy.

Acknowledgments

This study was supported by the Ministry of Science and Technology [105-2320-B-002-031-MY3, 105-2628-B-002-003-MY3].

Conflict of interest

The authors declare no conflict of interest.

References

Abraha, A.M., and Ketema, E.B. (2016). Apoptotic pathways as a therapeutic target for colorectal cancer treatment. *World J Gastrointest.*

- Oncol. 8: 583–591.
- Charoensinphon, N., Qiu, P., Dong, P., Zheng, J., Ngauv, P., Cao, Y., Li, S., Ho, C.T., and Xiao, H. (2013). 5-demethyltangeretin inhibits human nonsmall cell lung cancer cell growth by inducing G2/M cell cycle arrest and apoptosis. *Mol. Nutr. Food Res.* 57: 2103–2111.
- Chen, C., Ono, M., Takeshima, M., and Nakano, S. (2014). Antiproliferative and apoptosis-inducing activity of nobiletin against three subtypes of human breast cancer cell lines. *Anticancer Res.* 34: 1785–1792.
- Chen, Y.K., Wang, H.C., Ho, C.T., Chen, H.Y., Li, S., Chan, H.L., Chung, T.W., Tan, K.T., Li, Y.R., and Lin, C.C. (2015). 5-demethylnobiletin promotes the formation of polymerized tubulin, leads to G2/M phase arrest and induces autophagy via JNK activation in human lung cancer cells. *J Nutr. Biochem.* 26: 484–504.
- Chiou, Y.S., Sang, S., Cheng, K.H., Ho, C.T., Wang, Y.J., and Pan, M.H. (2013). Peracetylated (-)-epigallocatechin-3-gallate (ACEGCG) potently prevents skin carcinogenesis by suppressing the PKD1-dependent signaling pathway in CD34+ skin stem cells and skin tumors. *Carcinogenesis* 34: 1315–1322.
- Chiu, S.P., Wu, M.J., Chen, P.Y., Ho, Y.R., Tai, M.H., Ho, C.T., and Yen, J.H. (2013). Neurotrophic action of 5-hydroxylated polymethoxyflavones: 5-demethylnobiletin and gardenin A stimulate neurogenesis in PC12 cells. *J Agric. Food Chem.* 61: 9453–9463.
- Denlinger, C.S., and Barsevick, A.M. (2009). The challenges of colorectal cancer survivorship. *J Natl. Compr. Canc. Netw.* 7: 883–893.
- Hu, T., Li, Z., Gao, C.Y., and Cho, C.H. (2016). Mechanisms of drug resistance in colon cancer and its therapeutic strategies. *World J Gastroenterol.* 22: 6876–6889.
- Jemal, A., Siegel, R., Xu, J., and Ward, E. (2010). Cancer statistics, 2010. *CA Cancer J Clin.* 60: 277–300.
- Kawabata, K., Murakami, A., and Ohigashi, H. (2005). Nobiletin, a citrus flavonoid, down-regulates matrix metalloproteinase-7 (matrilysin) expression in HT-29 human colorectal cancer cells. *Biosci. Biotechnol. Biochem.* 69: 307–314.
- Kimura, J., Nemoto, K., Yokosuka, A., Mimaki, Y., Degawa, M., and Ohizumi, Y. (2013). 6-demethoxynobiletin, a nobiletin-analog citrus flavonoid, enhances extracellular signal-regulated kinase phosphorylation in PC12D cells. *Biol. Pharm. Bull.* 36: 1646–1649.
- Lai, C.S., Tsai, M.L., Cheng, A.C., Li, S., Lo, C.Y., Wang, Y., Xiao, H., Ho, C.T., Wang, Y.J., and Pan, M.H. (2011). Chemoprevention of colonic tumorigenesis by dietary hydroxylated polymethoxyflavones in azoxymethane-treated mice. *Mol. Nutr. Food Res.* 55: 278–290.
- Li, S., Lo, C.Y., and Ho, C.T. (2006). Hydroxylated polymethoxyflavones and methylated flavonoids in sweet orange (*Citrus sinensis*) peel. *J Agric. Food Chem.* 54: 4176–4185.
- Li, Y.H., Niu, Y.B., Sun, Y., Zhang, F., Liu, C.X., Fan, L., and Mei, Q.B. (2015). Role of phytochemicals in colorectal cancer prevention. *World J Gastroenterol.* 21: 9262–9272.
- Luo, G., Guan, X., and Zhou, L. (2008). Apoptotic effect of citrus fruit extract nobiletin on lung cancer cell line A549 in vitro and in vivo. *Cancer Biol. Ther.* 7: 966–973.
- Ma, N., Lai, C.S., Chung, C.H., Yang, J.M., Hsu, K.C., Chen, C.Y., Chun, T.S., Li, S., Ho, C.T., and Pan, M.H. (2014). 5-Demethyltangeretin is more potent than tangeretin in inhibiting dimethylbenz(a)anthracene (DMBA)/12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin tumorigenesis. *J Funct Foods* 11: 528–537.
- Moon, J.Y., Cho, M., Ahn, K.S., and Cho, S.K. (2013). Nobiletin induces apoptosis and potentiates the effects of the anticancer drug 5-fluorouracil in p53-mutated SNU-16 human gastric cancer cells. *Nutr. Cancer* 65: 286–295.
- Moon, J.Y., and Cho, S.K. (2016). Nobiletin Induces Protective Autophagy Accompanied by ER-Stress Mediated Apoptosis in Human Gastric Cancer SNU-16 Cells. *Molecules* 21(7): 914.
- Morley, K.L., Ferguson, P.J., and Koropatnick, J. (2007). Tangeretin and nobiletin induce G1 cell cycle arrest but not apoptosis in human breast and colon cancer cells. *Cancer Lett.* 251: 168–178.
- Nikoletopoulou, V., Markaki, M., Palikaras, K., and Tavernarakis, N. (2013). Crosstalk between apoptosis, necrosis and autophagy. *Biochim. Biophys. Acta* 1833: 3448–3459.
- Ouyang, L., Shi, Z., Zhao, S., Wang, F.T., Zhou, T.T., Liu, B., and Bao, J.K. (2012). Programmed cell death pathways in cancer: a review of apoptosis, autophagy and programmed necrosis. *Cell Prolif.* 45: 487–498.
- Pan, M.H., Chiou, Y.S., Cheng, A.C., Bai, N., Lo, C.Y., Tan, D., and Ho, C.T. (2007). Involvement of MAPK, Bcl-2 family, cytochrome c, and caspases in induction of apoptosis by 1,6-O-diacetylbritannilactone in human leukemia cells. *Mol. Nutr. Food Res.* 51: 229–238.
- Qiu, P., Dong, P., Guan, H., Li, S., Ho, C.T., Pan, M.H., McClements, D.J., and Xiao, H. (2010). Inhibitory effects of 5-hydroxy polymethoxyflavones on colon cancer cells. *Mol. Nutr. Food Res.* 54(2): S244–S252.
- Sergeev, I.N., Ho, C.T., Li, S., Colby, J., and Dushenkov, S. (2007). Apoptosis-inducing activity of hydroxylated polymethoxyflavones and polymethoxyflavones from orange peel in human breast cancer cells. *Mol. Nutr. Food Res.* 51: 1478–1484.
- Simon, K. (2016). Colorectal cancer development and advances in screening. *Clin. Interv. Aging* 11: 967–976.
- Song, M., Charoensinphon, N., Wu, X., Zheng, J., Gao, Z., Xu, F., Wang, M., and Xiao, H. (2016). Inhibitory Effects of Metabolites of 5-Demethylnobiletin on Human Nonsmall Cell Lung Cancer Cells. *J Agric. Food Chem.* 64: 4943–4949.
- Song, M., Wu, X., Charoensinphon, N., Wang, M., Zheng, J., Gao, Z., Xu, F., Li, Z., Li, F., Zhou, J., and Xiao, H. (2017). Dietary 5-demethylnobiletin inhibits cigarette carcinogen NNN-induced lung tumorigenesis in mice. *Food Funct.* 8: 954–963.
- Sp, N., Kang, D.Y., Joung, Y.H., Park, J.H., Kim, W.S., Lee, H.K., Song, K.D., Park, Y.M., and Yang, Y.M. (2017). Nobiletin Inhibits Angiogenesis by Regulating Src/FAK/STAT3-Mediated Signaling through PXN in ER(+) Breast Cancer Cells. *Int. J Mol. Sci.* 18(5): 935.
- Tuan, J., and Chen, Y.X. (2016). Dietary and Lifestyle Factors Associated with Colorectal Cancer Risk and Interactions with Microbiota: Fiber, Red or Processed Meat and Alcoholic Drinks. *Gastrointest. Tumors.* 3: 17–24.
- Tung, Y.C., Li, S., Huang, Q., Hung, W.L., Ho, C.T., Wei, G.J., and Pan, M.H. (2016). 5-Demethylnobiletin and 5-Acetoxy-6,7,8,3',4'-pentamethoxyflavone Suppress Lipid Accumulation by Activating the LKB1-AMPK Pathway in 3T3-L1 Preadipocytes and High Fat Diet-Fed C57BL/6 Mice. *J Agric. Food Chem.* 64: 3196–3205.
- Uckoo, R.M., Jayaprakasha, G.K., Vikram, A., and Patil, B.S. (2015). Polymethoxyflavones Isolated from the Peel of Miaray Mandarin (*Citrus miraya*) Have Biofilm Inhibitory Activity in *Vibrio harveyi*. *J Agric. Food Chem.* 63: 7180–7189.
- Wang, D., Wang, J., Huang, X., Tu, Y., and Ni, K. (2007). Identification of polymethoxylated flavones from green tangerine peel (*Pericarpium Citri Reticulatae Viride*) by chromatographic and spectroscopic techniques. *J Pharm. Biomed. Anal.* 44: 63–69.
- Wu, X., Song, M., Qiu, P., Li, F., Wang, M., Zheng, J., Wang, Q., Xu, F., and Xiao, H. (2018). A metabolite of nobiletin, 4'-demethylnobiletin and atorvastatin synergistically inhibits human colon cancer cell growth by inducing G0/G1 cell cycle arrest and apoptosis. *Food Funct.* 9: 87–95.
- Wu, X., Song, M., Wang, M., Zheng, J., Gao, Z., Xu, F., Zhang, G., and Xiao, H. (2015). Chemopreventive effects of nobiletin and its colonic metabolites on colon carcinogenesis. *Mol. Nutr. Food Res.* 59: 2383–2394.
- Yen, J.H., Weng, C.Y., Li, S., Lo, Y.H., Pan, M.H., Fu, S.H., Ho, C.T., and Wu, M.J. (2011). Citrus flavonoid 5-demethylnobiletin suppresses scavenger receptor expression in THP-1 cells and alters lipid homeostasis in HepG2 liver cells. *Mol. Nutr. Food Res.* 55: 733–748.
- Yoshimizu, N., Otani, Y., Saikawa, Y., Kubota, T., Yoshida, M., Furukawa, T., Kumai, K., Kameyama, K., Fujii, M., Yano, M., Sato, T., Ito, A., and Kitajima, M. (2004). Anti-tumour effects of nobiletin, a citrus flavonoid, on gastric cancer include: antiproliferative effects, induction of apoptosis and cell cycle deregulation. *Aliment. Pharmacol. Ther.* 20(1): 95–101.
- Zheng, J., Song, M., Dong, P., Qiu, P., Guo, S., Zhong, Z., Li, S., Ho, C.T., and Xiao, H. (2013a). Identification of novel bioactive metabolites of 5-demethylnobiletin in mice. *Mol. Nutr. Food Res.* 57: 1999–2007.